

MULTI-SATELLITE EARTH SCIENCE DATA RECORD FOR STUDYING GLOBAL VEGETATION TRENDS AND CHANGES

Kamel Didan

didan@email.arizona.edu

The University of Arizona, Tucson, AZ

Abstract—One of the stated goals of NASA Making Earth Science Data Records for Use in Research Environments ([MEaSUREs](http://measures.arizona.edu)) program is the support of the Earth Science research community by providing reliable Earth Science Data Records (ESDR). These products are expected not only to be of high quality but should also combine data from multiple sources to form the long and coherent measurements required for studying climate change impact on the Earth system.

To that end, this MEASUREs' project aims at generating a seamless and consistent sensor independent ESDR quality record of land surface vegetation index and phenology by fusing measurements from different satellite missions and sensors. We're using the AVHRR, MODIS, and VIIRS daily surface reflectance measurements, and the concept of homogeneous vegetation cluster model to design continuity algorithms that will be applied to these multi-sensor data sets. This effort will generate, characterize, and deliver 30+ years of consistent daily measurements of land surface VI and annual phenology parameters at a climate modeling grid resolution (0.05°, 5.6km). The consistency and accuracy of these products will be evaluated by comparison with in situ vegetation growing season observations over different biomes, latitudinal and elevational gradients. These ESDR products will be distributed through the USGS LP-DAAC. Key science and modeling community users, as well as the US National Phenology Network (US-NPN) will help with evaluating these ESDR products. Information about the project and the ESDR products status are available at <http://measures.arizona.edu>.

Keywords: *Remote Sensing, Vegetation Index, Land Surface Phenology, AVHRR, MODIS, climate change.*

I. INTRODUCTION

Vegetation indices (VI) from remote sensing are by far the most widely used remote sensing tools for studying vegetation and large-scale ecosystem processes [1]. In this context, knowledge of phenologic variability and the environmental conditions controlling their activity are further prerequisite to inter-annual studies and predictive modeling of land surface responses to climate change. Satellite phenology encompasses the analysis of the timing and rates of vegetation growth, senescence, and dormancy at seasonal and interannual time scales. Vegetation indices, which capture the aggregate functioning of a canopy, are the most robust and widely used measurement for extracting phenology information. Time series measures of MODIS enhanced vegetation index (EVI) have been shown highly correlated with flux tower photosynthesis in both tropical and temperate ecosystems at seasonal scales [2]. Changes in vegetation phenology depict an integrated response to change in environmental factors and provide valuable information to global change research.


Remote sensing of vegetation dynamic is based on the analysis of vegetation index time series profiles [3], because of their simplicity, stability, and inherent resistance to noise. Most vegetation index based studies are, however, limited to using one sensor owing to the inter-sensor continuity challenges. Although, phenology is used for a variety of research and application topics, the central premise remains the study of vegetation dynamics, changes and trends in response to change in climate and other factors. Consequently, the consistency and length of data records are key requirements. With satellite missions lasting only few years, long term studies of vegetation dynamics and phenology will have to combine multiple satellite data sources.

In the first phase of this project, our objective was to design a set of continuity algorithms to enable the seamless conversion of data across the different. This seamless record of land surface vegetation will be used by our land surface Phenology algorithm to estimate the timing and rates of vegetation change at seasonal and interannual time scales. Herein, we're reporting on our progress with this objective.

II. DATA AND METHOD

Although, the project will eventually use data from the AVHRR, MODIS, VIIRS [4], at this stage we're only working with the AVHRR and MODIS records (Table 1), since the VIIRS sensor is yet to fly. In order to facilitate the continuity work we plan to bridge the two records with SPOT-Vegetation sensor data from the period 1999 to 2002.

We're using daily global 0.05° resolution surface reflectance (AVH09C1) from 1981 to 1999 Advanced Very High Resolution Radiometer (AVHRR) Long Term Data Record (LTDR) [5] and daily Terra (available starting March 2000) and Aqua (available starting June 2002) MODerate Resolution Imaging Spectroradiometer (MODIS) Climate Modeling Grid daily surface reflectance (MOD09C1 and MYD09C1) [6]. Both records provide data in the red and near infrared (NIR) with different spectral characteristics (Figure 1), which is the primary cause of the discontinuity in these records.



Recent cross-sensor studies have shown the feasibility of NDVI and EVI translation across several sensor systems [7][8]. While NDVI requires only the Red and NIR bands, EVI extension, however, is limited to only sensors that carry a blue channel. To generate a backward compatible EVI data record that spans both AVHRR and MODIS we're implementing a 2-band adaptation of EVI, or EVI2 (Eq. 2), by eliminating the blue band which does not provide additional biophysical information, and was aimed at reducing noise and uncertainties associated with highly variable atmospheric aerosols [9].

$$EVI2 = 2.5 \frac{N-R}{N+2.4R+1} \quad (1)$$

To extend these measurements across AVHRR and MODIS sensor, we've developed a data fusion technique based on the homogeneous vegetation phenology clusters [10]. This method assumes that land surface vegetation behavior is the sum results of the controls exerted by climate, soil, elevation gradient, aspect, biological limitations and geographic location. The concept of phenology cluster is very similar to the biotic life-zones used to classify ecosystems or climate classification [11]. In that regard, a homogeneous vegetation phenology cluster is also a biotic zone, with similar plant species, at around the same elevation, and governed by similar temperature, precipitation and radiation regimes. These

clusters (Fig. 2) were constructed using elevation data (GTOPO30), mean annual temperature and precipitation [12], and the MODIS land cover map [13]. Inter-sensor-continuity is then modeled separately in each of these homogeneous clusters.

We computed a geometric regression [14] model that links the AVHRR LTDR with MODIS daily surface reflectance. The advantage of using a geometric model, as opposed to using the more common ordinary least square linear regression, is that it factors errors in both records. The regression model is then applied to the full record in each cluster enabling the conversion between the two sensors, and thus the creation of a sensor independent dataset.

III. RESULTS AND CONCLUSIONS

Our preliminary work indicates an overall high degree of agreement between the two records, owing to the VI ratioing capabilities. However, considerable differences (up to 30%) existed over the tropics and high latitude regions in the vegetation index signal. A large portion of this divergence resulted from poor data quality (aerosols, clouds, etc...). Using the data quality information we eliminated most poor quality data, and then applied the cluster based geometric models to the rest of the data.

The average difference between the two records was reduced to about 5% globally, well within current single sensor margin of error, insuring reliable data for global long term trends and change analysis. These early results support our hypothesis that a consistent and reliable sensor independent record is achievable using our cluster based continuity method.

Well calibrated, long term time series of vegetation indices and land surface phenology have wide social, cultural and economic significance to the well being of our planet, i.e., food supply, human health, invasive species, droughts, etc... In this work we're addressing a number of challenges, cross-discipline topics integrating multi-tiered phenology user communities, remote sensing, and modeling communities to advance our understanding of the Earth's terrestrial surface and biomes and their sustainability in the face of increasing human and environmental pressures. These records are critical to understanding Earth System processes, to assessing change and long term trends, provide input and validation means to modeling efforts, and support policy decisions and operational services.




Figure 2: Homogeneous vegetation phenology clusters map. Each of the unit represents an area with similar vegetation dynamic.

REFERENCES

- [1] K. Didan, A. Huete, A.R., "Analysis of the global vegetation dynamic metrics using MODIS Vegetation Index and land cover products," *Geoscience and Remote Sensing Symposium, 2004. IGARSS '04. Proceedings. 2004 IEEE International*, doi 10.1109/IGARSS.2004.1370755. 2004.
- [2] A.R. Huete, N. Restrepo-Coupe, P. Ratana, K. Didan, S.R. Saleska, K. Ichii, S. Panuthai, M. Gamo, "Multiple site tower flux and remote sensing comparisons of tropical forest dynamics in monsoon Asia," *Agric. & Forest Meteorology*, doi 10.1016/j.agrformet.2008.01.012. 2008.
- [3] X. Zhang, MA. Friedl, CB. Schaaf, AH. Strahler, JCF. Hodges, F. Gao, *et al.*, "Monitoring vegetation phenology using MODIS," *Remote Sensing of Environment*, **84** (2003), pp. 471–475. 2003.

- [4] R.E. Murphy, "The NPOESS preparatory project. Earth Science Satellite Remote Sensing," *Springer Berlin Heidelberg* (2006), pp. 182–198 [10.1007/978-3-540-37293-6](https://doi.org/10.1007/978-3-540-37293-6). 2006.
- [5] J. Pedelty, S. Devadiga, E.B. Masuoka, J.M. Pinzon, C.J. Tucker and E.F. Vermote *et al.*, "Generating a long-term land data record from the AVHRR and MODIS Instruments," *Geoscience and Remote Sensing Symposium, 2007, IGARSS 2007. IEEE International (2007)*, pp. 1021–1025. 2007.
- [6] C. Justice, D. Hall, V. Salomonson, J. Privette, G. Riggs, G., Strahler, et al., "The Moderate Resolution Imaging Spectroradiometer (MODIS): Land remote sensing for global change research," *IEEE Trans. Geosci. and Remote Sens.*, 36(4):1228-1249. 1998.
- [7] K. Gallo K, L. Ji, B. Reed, J. Idenshink, J. Dwyer, "Multi-platform comparisons of MODIS and AVHRR normalized Difference Vegetation Index data," *Remote Sensing of Environment*, 99, 221–231. 2005.
- [8] T. Miura, A.R. Huete, H. Yoshioka, "An empirical investigation of cross-sensor relationships of NDVI and red/near-infrared reflectance using EO-1 hyperion data," *Remote Sensing of Environment*, 100, 223-236. 2006.
- [9] J. Zhangyan, A.R. Huete, K. Didan, T. Miura, "Development of a two-band enhanced vegetation index without a blue band," *Remote Sensing of Environment* **112** (2008), pp. 3833-3845. doi:10.1016/j.rse.2008.06.006. 2008.
- [10] M.A. White, K.M. de Beurs, K. Didan, D.W. Inouye, et al., "Intercomparison, interpretation, and assessment of spring phenology in North America estimated from remote sensing for 1982 to 2006," *Global Change Biology*, 15 2335-59. 2009.
- [11] M. Kottek, J. Grieser, C. Beck, B. Rudolf, F. Rubel, "World map of the Köppen-Geiger climate classification updated, Meteorol. Zeitschr., 15(3), 259–263. 2006.
- [12] R. Leemans, W. Cramer, "The IIASA database for mean monthly values of temperature, precipitation and cloudiness on a global terrestrial grid", *Research Report RR-91-18. November 1991. International Institute of Applied Systems Analyses, Laxenburg*, pp. 61. 1991.
- [13] M.A. Friedl, D.K. McIver, J.C.F. Hodges, X.Y. Zhang, et al., "Global land cover mapping from MODIS: algorithms and early results", *Remote Sensing of Environment*, 83, 287– 302. 2002.
- [14] L. Leng, T. Zhang L. Kleinmann, W. Zhu W, "Ordinary least square regression, orthogonal regression, geometric mean regression and their applications in aerosol science", *Journal of Physics: Conference Series* **78**: 1-5. DOI. [10.1088/1742-6596/78/1/012084](https://doi.org/10.1088/1742-6596/78/1/012084). 2007.